

# PHYSICS 1A

## Quiz #3A

H. E. Smith

7 May 2004

*Closed book and notes; only these sheets, Scantron, pen/pencil and calculator may be used. This is exam version A. Mark your Scantron Test Form "A" and code your UCSD ID# under ID Number. Write your Name and ID number on the second sheet of this Quiz. Answer Questions 1-5 on the Scantron; answer Problem 6 in the space provided. Organize your work if you want to be considered for partial credit. GOOD LUCK!*

Useful Formulae:

$$\text{net}\vec{F} = m\vec{a}$$

$$F_{grav} = mg$$

$$f_s \leq \mu_s N \quad f_k = \mu_k N$$

$$\langle \vec{v} \rangle = \frac{\Delta \vec{x}}{\Delta t} \quad \vec{v} = \frac{d\vec{x}}{dt}$$

$$\langle \vec{a} \rangle = \frac{\Delta \vec{v}}{\Delta t} \quad \vec{a} = \frac{d\vec{v}}{dt} = \frac{d^2 \vec{s}}{dt^2}$$

$$x = x_0 + v_{0x} \cdot t + \frac{1}{2} a_x \cdot t^2 \quad y = y_0 + v_{0y} \cdot t + \frac{1}{2} a_y \cdot t^2$$

$$v_x = v_{0x} + a_x t \quad v_y = v_{0y} + a_y t$$

$$g = -9.80 m s^{-2}$$

$$W = \int F_{\parallel} ds$$

$$PE_{grav} = mgh \quad KE = \frac{1}{2} m v^2$$

$$\text{Quadratic : } ax^2 + bx + c = 0 \quad \rightarrow \quad x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

1. A student lifts a 20 kg box to a height of 1.5m, then walks a distance of 10m. The work done by the student is about
  - a) 200J
  - b) 230J
  - c) 300J
  - d) 2000J
  - e) 2300J
  
2. You push a 100kg crate 10m across the floor with a horizontal force of 200N. The coefficients of friction between the crate and the floor are  $\mu_s = 0.2$  and  $\mu_k = 0.1$ . If the crate starts at rest, the final speed is about
  - a) 2 m/s
  - b) 4.5 m/s
  - c) 10 m/s
  - d) 20 m/s
  - e) 22.5 m/s.
  
3. A car traveling at  $v = 10\text{m/s}$  skids to a stop in a distance of 10m. If the same car is traveling with  $v = 20\text{m/s}$ , how far will it skid before stopping?
  - a) 14m.
  - b) 20m.
  - c) 28m.
  - d) 40m.
  - e) 100m.
  
4. A mass,  $M$ , is attached to a massless rod with length,  $L$ , to act as a pendulum. The mass is lifted through a  $90^\circ$  angle so that the rod is horizontal, then released. At the bottom of its swing the pendulum has velocity
  - a)  $v = (gL)^2$
  - b)  $v = gL$
  - c)  $v = \sqrt{2gL}$
  - d)  $v = MgL$
  - e)  $v = \sqrt{MgL}$
  
5. Two Physics 1 students of equal mass sit at the top of a playground slide which makes an angle of  $45^\circ$  with the ground. Student #1 falls off the slide (vertically downward) as student #2 starts sliding downward. Assuming that they both start off with  $KE = 0$ , then, when they reach the ground
  - a)  $KE_1 = 2KE_2$
  - b)  $KE_1 = \sqrt{2}KE_2$
  - c)  $KE_1 = KE_2$
  - d)  $KE_1 = \frac{1}{\sqrt{2}}KE_2$
  - e)  $KE_1 = \frac{1}{2}KE_2$

Name \_\_\_\_\_

ID # A \_\_\_\_\_

6) (13 pts) A sled with mass  $m = 50\text{kg}$  slides up a hill at  $30^\circ$  inclination and then back down as shown. The velocity of the sled as it starts up the hill is  $5\text{m/s}$  and it experiences a frictional force of  $50\text{N}$  both on its way up and on its way down. How far up the hill does the sled slide?

 $d =$  \_\_\_\_\_

In the spaces below draw graphs of the KE, PE and total mechanical energy of the sled as a function of time.